



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

## BRAZIL.

*Sanitary report from Rio.*RIO DE JANEIRO, *January 3, 1899.*

SIR: I have the honor to transmit to you the official report for the week ended January 20. There were 283 deaths from all causes, a decrease of 15 as compared with the foregoing week; 10 deaths from *accessio pernicioso*, a decrease of 5; 13 deaths from yellow fever, a decrease of 7; 5 deaths from smallpox, a decrease of 2; 3 deaths from typhoid fever, none before; 1 death from diphtheria, none before; 1 death from measles, a decrease of 1; 2 deaths from beriberi, the same as before, and 68 deaths from tuberculosis, and increase of 16.

From the state of Sao Paulo I can give you some official data concerning the sanitary conditions during the month of November. The total number of deaths from all causes was 2,431, and increase of 1,179, as compared with the month of October, and from tuberculosis there died 142 persons, a decrease of 17, and from malarial fever, 71 persons, an increase of 13.

From typhoid fever there died 28 persons (11 at the city of Sao Paulo), an increase of 2; from yellow fever none, and from smallpox 20, an increase of 5; 15 cases occurred at the city of Sao Paulo, 1 at Pantos, and 4 at Una.

In regard to other infectious diseases, there were in the state of Sao Paulo 5 deaths from measles, 3 deaths from diphtheria, 36 deaths from whooping cough, 2 deaths from erysipelas, 17 deaths from influenza, 18 deaths from dysentery, 2 deaths from leprosy, 7 deaths from cholera nostras, and 1 death from beriberi.

Since last report the following-named ships have been inspected and received bills of health of this office: January 30, steamship *Grecian Prince*, British, for New York; steamship *Bellarden*, British, for New York. February 1, steamship *Ivydene*, British, for New York.

Respectfully, yours,

W. HAVELBURG, M. D.,

*Sanitary Inspector, U. S. M. H. S.*

The SUPERVISING SURGEON-GENERAL,

*U. S. Marine-Hospital Service.*

## CUBA.

*Sanitary report from Havana.*HAVANA, CUBA, *March 3, 1899.*

SIR: I have the honor to submit the following report for the week ended Thursday, March 2, 1899:

In all probability the death rate during this week has been the lowest recorded since 1894. There have been but 187 deaths during this time from all causes. No deaths are reported either from yellow fever or smallpox.

There seems to be an increasing death rate from diphtheria, 5 deaths from that disease having occurred during the week.

A slight increase is also shown in deaths from different malarial fevers.

Several cases of suspicious fever have been reported among the regulars stationed in the city and in the volunteer forces of the Seventh Army Corps in and around Marianno. From the medical staff of the Army I learn that none of these cases have been designated as yellow fever.